

REFERENCIAS BIBLIOGRAFICAS

- 1.- Harris, MI; Klein, R; Welborn, TA and Knuiiman , MW .**Association Onset of NIDDM occurs at least 4-7 yr before clinical diagnosis.** Diabetes Care 1992; 15(7): 815-819.
- 2.- Klein, BE; Klein, R; Moss, SE and Cruickshanks, KJ. **Parental history of diabetes in a population-based study.** Diabetes Care 1996; 19(8): 827-830.
- 3.-Thorand, Barbara; Liese, Angela D; Metzger, Marie-Hélène; Reitmeir, Peter; Schneider, Andrea and Löwel, Hannelore .**Can inaccuracy of reported parental history of diabetes explain the maternal transmission hypothesis for diabetes?** International Journal of Epidemiology 2001;30:1084-1089.
- 4.- Meigs James B., Cupples L. Adrienne, and Wilson Peter W.F. **Parental Transmission of Type 2 Diabetes. The Framingham Offspring Study.** Diabetes 2000; 49:2201-2207.
- 5.- Kohner, Eva M.; Aldington, Stephen J. ; Stratton, Irene M. ; Manley, Susan E. ; Holman, Rury R. ; Matthews, David R. et al. **United Kingdom Prospective Diabetes Study Diabetic Retinopathy at Diagnosis of Non-Insulin-Dependent Diabetes Mellitus and Associated Risk Factors.** Arch Ophthalmol. 1998;116:297-303.
- 6.- Larson, TS ; Santanello, N; Shahinfar, S; O'Brien, PC; Palumbo, PJ ; Melton, LJ et al . **Association Trends in persistent proteinuria in adult-onset diabetes: a population-based study.** Diabetes Care 2000; 23(1): 51-56.
- 7.- Sosenko, Jay M. ; Hu, Dongsheng; Welty, Tom; Howard, Barbara V. ; Lee, Elisa and Robbins, David C. **Albuminuria in Recent-Onset Type 2 Diabetes The Strong Heart Study.** Diabetes Care 2002; 25:1078-1084.
8. - Hillier ,Teresa A. ; Pedula, Kathryn L. **Characteristics of an Adult Population With Newly Diagnosed Type 2 Diabetes The relation of obesity and age of onset.** Diabetes Care 2001; 24:1522-1527..
- 9.- Wens, Johan; Van Casteren, Viviane; Vermeire, Etienne; Van Royen, Paul; Denekens, Joke. **Newly Diagnosed Type 2 Diabetic Population in Belgium** Diabetes Care 2002; 25:637-638.
- 10.- Spijkerman, Annemieke M.W. ; Adriaanse, Marcel C. ; Dekker, Jacqueline M. ; Nijpels, Giel ; Stehouwer, Coen D.A. ; Bouter, Lex M. et al. **Diabetic**

Patients Detected by Population-Based Stepwise Screening Already Have a Diabetic Cardiovascular Risk Profile Diabetes Care 2002; 25:1784-1789.

11.- UK Prospective Diabetes Study Group. **Control intensivo con sulfonilureas o insulina en comparación con un tratamiento convencional y riesgo de complicaciones en pacientes con diabetes tipo 2 (UKPDS 33)** The Lancet 33(5): 258-287 1998.

12.-Gamarra M., Alvizuri H. **Características clínicas y epidemiológicas del debut de diabetes mellitus tipo 2 en el Hospital Antonio Lorena del Cusco.** IX Congreso Peruano de Endocrinología 2002.

13.- Buchanan, Thomas A. **Pancreatic B-Cell Defects in Gestational Diabetes: Implications for the Pathogenesis and Prevention of Type 2 Diabetes** .The Journal of Clinical Endocrinology & Metabolism 86(3): 989-993 2001.

14.- Léo Wajchenberg, Bernardo. **Subcutaneous and Visceral Adipose Tissue: Their Relation to the Metabolic Syndrome.** Endocrine Reviews 21 (6): 697-738 2000.

15.- American Diabetes Association. **Management of Dyslipidemia in Adults With Diabetes.** Diabetes Care 25:S74-S77, 2002.

16.- Henry , Robert R. **Preventing Cardiovascular Complications of Type 2 Diabetes: Focus on Lipid Management.** Clinical Diabetes 19(3): 113-120 2001.

17.- Lauenborg, J.; Hansen ,T.; Møller Jensen, D.; Vestergaard ,H.; Mølsted Pedersen,L.; Hornnes ,P. , et al . **Increasing Incidence of Diabetes After Gestational Diabetes. A long-term follow-up in a Danish population** Diabetes Care 27:1194-1199, 2004.