

VII. BIBLIOGRAFIA

1. Schaumburg-Lever G, Safford OE, Orfanos CE, Lever WF: Herpes gestationis: histology and ultrastructure. Arch Dermatol 107: 888-892, 1973.
2. Shornick JK, Bangert JL, Freeman RG, Gilliam JN: *Herpes gestationis*: clinical and histologic features of twenty-eight cases. J Am Acad Dermatol 8: 214-24, 1983.
3. Jordon RE, Heine KG, Tappeiner G et al.: The immunopathology of *Herpes gestationis*. Immunofluorescence studies and characterization of 'HG factor'. J Clin Invest 57: 1426-1431, 1976.
4. Katz SI, Hertz K, Yaoita H: *Herpes gestationis*. Immunopathology and characterization of the HG factor. J Clin Invest 57: 1434-1441, 1976.
5. Morrison LH, Labib RS, Zone JJ, Diaz LA, Anhalt GJ: *Herpes gestationis* autoantibodies recognize a 180-kD human epidermal antigen. J Clin Invest 87: 2023-2026, 1988.
6. Kelly SE, Bhogal BS, Wojnarowska F, Whitehead P, Leigh IM, Black MM: Westem blot analysis of the antigen in pemphigoid gestationis. Br J Dermatol 122: 445- 449, 1990.
7. Giudice GJ, Emery DJ, Zelickson BD, Anhalt GJ, Liu Z, Diaz LA: Bullous pemphigoid and herpes gestationis autoantibodies recognize a common non-collagenous site on the BP180 ectodomain. J Immunol 151: 5742-50, 1993.
8. Giudice GJ, Wilske KC, Anhalt GJ, Fairley JA, Taylor AF, Emery DJ, Hoffman RG, Diaz LA: Development of an ELISA to detect anti-BP180 autoantibodies in bullous pemphigoid and herpes gestationis. J Invest Dermatol 102: 878-881,1994.

9. Hodge L, Black MM, Ramnarain N, Bhogal BS: Indirect complement immunofluorescence in the assessment of bullous pemphigoid, cicatricial pemphigoid and herpes gestationis. *Clin Exp Dermatol* 3: 61-67, 1978.
10. Gammon WR, Briggaman RA, Inman AO, Queen LL, Wheeler CE: Differentiating anti-lamina lucida and anti-sublamina densa anti-BMZ antibodies by indirect immunofluorescence on 1.0 M sodium chloride separated skin. *J Invest Dermatol* 82: 139-144, 1984.
11. Hashimoto T, Ogawa MM, Konohana A, Nishikawa T: Detection of pemphigus vulgaris and pemphigus foliaceus antigens by immunoblot analysis using different antigen sources. *J Invest Dermatol* 94: 327-331, 1990.
12. Sugi T, Hashimoto T, Hibi T, Nishikawa T. Production of human monoclonal anti-basement membrane zone (BMZ) antibodies from a patient with bullous pemphigoid (BP) by Epstein-Barr virus transformation. Analysis of the heterogeneity of anti-BMZ antibodies in BP sera using them. *J Clin Invest* 84: 1050-1055, 1989.
13. Giudice GJ, Emery DJ, Diaz LA. Cloning and primary structural analysis of the bullous pemphigoid autoantigen, BP-180. *J Invest Dermatol* 99: 243-250, 1992.
14. Amagai M, Hashimoto T, Tajima S, Inokuchi Y, Shimizu N, Saito M, Miki K, Nishikawa T: Partial cDNA cloning of the 230-kD mouse bullous pemphigoid antigen by use of a human monoclonal anti-basement membrane zone antibody. *J Invest Dermatol* 95: 252-259, 1990.
15. Tanaka M, Hashimoto T, Amagai M, Shimizu N, Ikegami N, Tsubata T, Hasegawa A, Miki K, Nishikawa T: Characterization of bullous pemphigoid antibodies by use of recombinant bullous pemphigoid antigen proteins. *J Invest Dermatol* 97: 725-728, 1991.

16. Shornick JK: Herpes gestationis. J Am Acad Dermatol 17: 539-556,1987.
17. Bushkell LL, Jordon RE, Goltz RW: Herpes gestationis: new immunologic findings. Arch Dermatol 110: 65-69, 1974.
18. Kirtschig G, Collier PM, Emmerson RW, Wojnarowska F: Severe case of pemphigoid gestationis with unusual target antigen. Br J Dermatol 131: 108-11, 1994.
19. Liu Z, Diaz LA, Troy JL, Taylor AF, Emery DJ, Fairley JA, Giudice GJ: A passive transfer model of the organ-specific autoimmune disease, bullous pemphigoid, using antibodies generated against the hemidesmosomal antigen, BP180. J Clin Invest 92: 2480-2488, 1993.